

25



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 253/30, 255/36, 255/37, 255/38, 255/41, 255/46, A61K 31/275</b>		A1	(11) International Publication Number: <b>WO 95/24381</b>
			(43) International Publication Date: 14 September 1995 (14.09.95)
(21) International Application Number: PCT/US95/02965 (22) International Filing Date: 8 March 1995 (08.03.95) (30) Priority Data: 9404706.5                      11 March 1994 (11.03.94)                      GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; Corporate, Intellectual Property, WW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BORDAS-NAGY, Joseph [CA/US]; Apartment 0102, 655 South Fair Oaks, Sunnyvale, CA 94086 (US). GORYCKI, Peter [US/US]; 1801 Butler Pike #121, Conshohocken, PA 19428 (US). WEBB, Kevin, Scott [US/US]; 1415 Powder Horn Drive, Phoenixville, PA 19460 (US). (74) Agents: KANAGY, James, M. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, PT, RO, RU, SD, SG, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published With international search report.	
(54) Title: CYANO COMPOUNDS AND PREPARATIONS THEREOF			
(57) Abstract			
<p>This invention relates to compounds of formula (I) which are useful as PDE IV inhibitors and for treating diseases related thereto.</p> <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>			

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**BEST AVAILABLE COPY**

## CYANO COMPOUNDS AND PREPARATIONS THEREOF

**Field of the Invention**

The present invention relates to novel phenylcyclohexan-1-ylcarboxylic acids, pharmaceutical compositions containing these compounds, the use of these compounds in  
5 treating allergic and inflammatory diseases and to the use of these compounds to inhibit the production of Tumor Necrosis Factor (TNF).

**Background of the Invention**

Bronchial asthma is a complex, multifactorial disease characterized by reversible narrowing of the airway and hyperreactivity of the respiratory tract to external stimuli.  
10 Identification of novel therapeutic agents for asthma is made difficult by the fact that multiple mediators are responsible for the development of the disease. Thus, it seems unlikely that eliminating the effects of a single mediator will have a substantial effect on all components of chronic bronchial asthma.

An alternative to the "mediator approach" is to regulate the activity of cells  
15 responsible for the pathophysiology of asthma. Cyclic AMP (cAMP, adenosine cyclic 3',5'-monophosphate) modulates the activity of most, if not all, of the cells that contribute to the pathophysiology of extrinsic (allergic) asthma. An elevation of cAMP would produce beneficial effects including: (1) airway smooth muscle relaxation, (2) inhibition of mast cell mediator release, (3) suppression of neutrophil degranulation, (4) inhibition of basophil  
20 degranulation, and (5) inhibition of monocyte and macrophage activation. Cyclic AMP has been shown to mediate cellular responses to a wide range of hormones, neurotransmitters and drugs; [Krebs Endocrinology Proceedings of the 4th International Congress Excerpta Medica, 17-29, 1973].

One potential means to regulate the activity of cells responsible for the  
25 pathophysiology of asthma is to control the intracellular levels of cyclic AMP. Cellular cAMP levels are elevated when an appropriate agonist binds to particular cell surface receptors, thereby activating adenylate cyclase to convert  $Mg^{+2}$ -ATP to cAMP at an accelerated rate. The principal cellular mechanism for the inactivation of cAMP is hydrolysis of the 3'-phosphodiester bond by one or more of a family of isozymes referred to  
30 as cyclic nucleotide phosphodiesterases (cyclic nucleotide phosphodiesterase hereinafter "PDE"s). Hence, compounds that activate adenylate cyclase or inhibit phosphodiesterase should be effective in suppressing the inappropriate activation of airway smooth muscle and a wide variety of inflammatory cells.

It has been shown that a distinct PDE isozyme, PDE IV, is responsible for cAMP  
35 breakdown in airway smooth muscle and inflammatory cells. [Torphy, "Phosphodiesterase Isozymes: Potential Targets for Novel Anti-asthmatic Agents" in New Drugs for Asthma, Barnes, ed. IBC Technical Services Ltd., 1989]. Research indicates that inhibition of this enzyme not only produces airway smooth muscle relaxation, but also suppresses

degranulation of mast cells, basophils and neutrophils along with inhibiting the activation of monocytes and neutrophils. The beneficial effects of PDE IV inhibition are markedly potentiated when adenylate cyclase activity of target cells is elevated by appropriate hormones or autocooids. Thus, PDE IV inhibitors would be effective in the asthmatic lung, where levels of prostaglandin E<sub>2</sub> and prostacyclin (both activators of adenylate cyclase) are elevated. PDE IV inhibitors offer a unique approach to the pharmacotherapy of bronchial asthma, and possess significant therapeutic advantages over agents currently on the market. The compounds of this invention have the ability to inhibit PDE IV.

The compounds of this invention also inhibit the production of TNF, a serum glycoprotein. Excessive or unregulated TNF production has been implicated in mediating or exacerbating a number of undesirable physiological conditions, such as diseases, and including rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions; sepsis, septic shock, endotoxic shock, gram negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, cerebral malaria, chronic pulmonary inflammatory disease, silicosis, pulmonary sarcoidosis, bone resorption diseases, reperfusion injury, graft vs. host reaction, allograft rejections, fever and myalgias due to infection, such as influenza, cachexia secondary to infection or malignancy, human acquired immune deficiency syndrome (AIDS), cachexia secondary to AIDS, AIDS related complex (ARC), keloid formation, scar tissue formation, Crohn's disease, ulcerative colitis, or pyresis, in addition to a number of autoimmune diseases, such as multiple sclerosis, autoimmune diabetes and systemic lupus erythematosus.

AIDS results from the infection of T lymphocytes with Human Immunodeficiency Virus (HIV). At least three types or strains of HIV have been identified: HIV-1, HIV-2 and HIV-3. As a consequence of HIV infection, T-cell-mediated immunity is impaired and infected individuals manifest severe opportunistic infections and/or unusual neoplasms. HIV entry into a T lymphocyte requires prior T lymphocyte activation. Once an activated T lymphocyte has been infected with HIV, the T lymphocyte must be maintained in an activated state in order to permit HIV gene expression and/or HIV replication.

Cytokines, including TNF, are implicated in activated T-cell-mediated HIV protein expression and/or virus replication as playing a role in maintaining T lymphocyte activation. Therefore, interference with cytokine activity in an HIV-infected individual, such as by inhibition of TNF production, aids in limiting the maintenance of T cell activation, and thereby mitigates the progression of HIV infection to previously uninfected cells. When HIV infection of previously uninfected cells is diminished, a slowing or elimination of the progression of immune dysfunction caused by HIV infection results.

Monocytes, macrophages, and related cells, such as kupffer and glial cells, have also been implicated in the maintenance of HIV infection. These cells, like T cells, are targets for viral replication, where the level of viral replication is dependent upon the


- activation state of the cells. [See Rosenberg *et al.*, The Immunopathogenesis of HIV Infection, Advances in Immunology, Vol. 57, 1989]. Monokines, such as TNF, have been shown to activate HIV replication in monocytes and/or macrophages [See Poli *et al.*, Proc. Natl. Acad. Sci., 87:782-784, 1990], therefore, inhibition of monokine production or
- 5 activity aids in limiting HIV progression as stated above for T cells.

TNF has also been implicated in various roles with other viral infections, such as the cytomegalovirus (CMV), influenza virus, adenovirus, and the herpes virus for similar reasons as those noted. TNF is also associated with yeast and fungal infections. Specifically *Candida albicans* has been shown to induce TNF production *in vitro* in human

10 monocytes and natural killer cells. [See Riipi *et al.*, Infection and Immunity, 58(9):2750-54, 1990; and Jafari *et al.*, Journal of Infectious Diseases, 164:389-95, 1991. See also Wasan *et al.*, Antimicrobial Agents and Chemotherapy, 35(10):2046-48, 1991; and Luke *et al.*, Journal of Infectious Diseases, 162:211-214, 1990].

### 15 Summary of the Invention

This invention relates to certain compounds of Formula I



20 wherein

R<sub>1</sub> is OH or an ether or ester thereof,

X is YR<sub>2</sub>, halogen, nitro, NR<sub>4</sub>R<sub>5</sub> or formyl amine;

Y is O or S(O)<sub>m</sub>, where m is 0, 1 or 2;

25 R<sub>2</sub> is methyl or ethyl, where either methyl or ethyl may be optionally substituted by 1 or more halogens;

R<sub>3</sub> is hydrogen, halogen, C<sub>1-4</sub>alkyl, CH<sub>2</sub>NHC(O)C(O)NH<sub>2</sub>, halo-substituted C<sub>1-4</sub>alkyl, -CH=CHR<sub>8</sub>R<sub>8</sub>', cyclopropyl optionally substituted by R<sub>8</sub>', CN, OR<sub>8</sub>, CH<sub>2</sub>OR<sub>8</sub>, NR<sub>8</sub>R<sub>10</sub>, CH<sub>2</sub>NR<sub>8</sub>R<sub>10</sub>, C(Z')H, C(O)OR<sub>8</sub>, C(O)NR<sub>8</sub>R<sub>10</sub>, or -C≡CR<sub>8</sub>';

R<sub>4</sub> and R<sub>5</sub> are independently hydrogen or C<sub>1-2</sub> alkyl;

30 R<sub>7</sub> is -(CR<sub>4</sub>R<sub>5</sub>)<sub>q</sub>R<sub>12</sub> or C<sub>1-6</sub> alkyl wherein the R<sub>12</sub> or C<sub>1-6</sub> alkyl group is optionally substituted one or more times by C<sub>1-2</sub> alkyl optionally substituted by one to three groups selected from -F, -Br, -Cl, -NO<sub>2</sub>, -NR<sub>10</sub>R<sub>11</sub>, -C(=O)R<sub>8</sub>, -C(=O)OR<sub>8</sub>, -OR<sub>8</sub>, -CN, -C(=O)NR<sub>10</sub>R<sub>11</sub>, -OC(=O)NR<sub>10</sub>R<sub>11</sub>, -OC(=O)R<sub>8</sub>, -NR<sub>10</sub>C(=O)NR<sub>10</sub>R<sub>11</sub>, -NR<sub>10</sub>C(=O)R<sub>11</sub>, -NR<sub>10</sub>C(=O)OR<sub>9</sub>, -NR<sub>10</sub>C(=O)R<sub>13</sub>, -C(=NR<sub>10</sub>)NR<sub>10</sub>R<sub>11</sub>, -C(=N-

CN)NR<sub>10</sub>R<sub>11</sub>, -C(=N-CN)SR<sub>9</sub>, -NR<sub>10</sub>C(=N-CN)NR<sub>10</sub>R<sub>11</sub>, -NR<sub>10</sub>S(=O)<sub>2</sub>R<sub>9</sub>,  
 -S(O)<sub>m</sub>R<sub>9</sub>, -NR<sub>10</sub>C(=O)C(=O)NR<sub>10</sub>R<sub>11</sub>, -NR<sub>10</sub>C(=O)C(=O)R<sub>10</sub>, thiazolyl, imidazolyl,  
 oxazolyl, pyrazolyl, triazolyl or tetrazolyl;

R<sub>8</sub> is -H or R<sub>9</sub>;

5 R<sub>8</sub>' is R<sub>8</sub> or fluorine;

R<sub>9</sub> is C<sub>1-4</sub> alkyl optionally substituted by one to three -F;

R<sub>10</sub> is OR<sub>8</sub>, hydrogen, or C<sub>1-4</sub>alkyl optionally substituted by one to three  
 fluorines;

10 R<sub>11</sub> is -H or C<sub>1-4</sub> alkyl optionally substituted by one to three -F; or when R<sub>10</sub> and  
 R<sub>11</sub> are as NR<sub>10</sub>R<sub>11</sub> they may together with the nitrogen form a 5 to 7 membered ring  
 optionally containing at least one additional heteroatom selected from O, N or S;

15 R<sub>12</sub> is C<sub>3-7</sub> cycloalkyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrimidyl, pyrazolyl,  
 1-imidazolyl, 2-imidazolyl, thiazolyl, triazolyl, pyrrolyl, piperaziny, piperidinyl,  
 morpholinyl, furanyl, 2-thienyl, 3-thienyl, 4-thiazolyl, 5-thiazolyl, quinolinyl, naphthyl or  
 phenyl;

20 R<sub>13</sub> is a heterocyclic ring selected from oxazolidinyl, oxazolyl, thiazolyl, pyrazolyl,  
 triazolyl, tetrazolyl, imidazolyl, imidazolidinyl, thiazolidinyl, isoxazolyl, oxadiazolyl or  
 thiadiazolyl, where R<sub>13</sub> is appended to a compound of Formula (I) through a carbon atom  
 of the heterocyclic ring, and where each heterocyclic ring may be unsubstituted or  
 substituted by one or two C<sub>1-2</sub> alkyl groups;

R<sub>14</sub> is H or R<sub>7</sub>, or when R<sub>10</sub> and R<sub>14</sub> are as NR<sub>10</sub>R<sub>14</sub>, they may together with the  
 nitrogen atom form a 5 to 7 membered ring optionally containing at least one additional  
 heteroatom selected from O, N or S;

m' is 0, 1 or 2;

25 q is 0, 1 or 2.

Y' is O or S;

30 Z is C(=Y')R<sub>14</sub>, C(=O)OR<sub>14</sub>, C(=Y')NR<sub>10</sub>R<sub>14</sub>, C(=NR<sub>10</sub>)NR<sub>10</sub>R<sub>14</sub>, CN,  
 C(=NOR<sub>8</sub>)R<sub>14</sub>, C(=O)NR<sub>8</sub>NR<sub>8</sub>C(=O)R<sub>8</sub>, C(=O)NR<sub>8</sub>NR<sub>10</sub>R<sub>14</sub>, C(=NOR<sub>14</sub>)R<sub>8</sub>,  
 C(=NR<sub>8</sub>)NR<sub>10</sub>R<sub>14</sub>, C(=NR<sub>14</sub>)NR<sub>8</sub>R<sub>8</sub>, C(=N-CN)NR<sub>10</sub>R<sub>14</sub>, C(=N-CN)SR<sub>9</sub>, 2-  
 imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 4-  
 triazolyl[1,2,3], 5-triazolyl[1,2,3], 3-triazolyl[1,2,4], 5-triazolyl[1,2,4], 5-tetrazolyl, 2-  
 oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-  
 oxadiazolyl[1,2,4], 5-oxadiazolyl[1,2,4], 2-oxadiazolyl[1,3,4], 2-thiadiazolyl[1,3,4], 2-  
 thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-oxazolidinyl, 4-oxazolidinyl, 5-oxazolidinyl, 2-  
 35 thiazolidinyl, 4-thiazolidinyl or 5-thiazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, or 5-  
 imidazolidinyl; wherein all of the heterocyclic ring systems may be optionally substituted  
 one or more times by R<sub>14</sub>;

Z' is O, NR<sub>9</sub>, NOR<sub>8</sub>, NCN, C(-CN)<sub>2</sub>, CR<sub>8</sub>CN, CR<sub>8</sub>NO<sub>2</sub>, CR<sub>8</sub>C(O)OR<sub>8</sub>, CR<sub>8</sub>C(O)NR<sub>8</sub>R<sub>8</sub>, C(-CN)NO<sub>2</sub>, C(-CN)C(O)OR<sub>9</sub>, or C(-CN)C(O)NR<sub>8</sub>R<sub>8</sub> ;

or

a pharmaceutically acceptable salt thereof.

5 This invention also relates to pharmaceutical compositions comprising a compound of Formula (I) and a pharmaceutically acceptable excipient.

This invention also relates to a method of mediation or inhibition of the enzymatic activity (or catalytic activity) of PDE IV in mammals, including humans, which comprises administering to a mammal in need thereof an effective amount of a compound of Formula  
10 (I) or a pharmaceutically acceptable salt thereof.

This invention also relates to a method for the treatment of allergic and inflammatory disease which comprises administering to a mammal, including humans, in need thereof, an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

15 This invention also relates to a method for the treatment of asthma which comprises administering to a mammal, including humans, in need thereof, an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

This invention further relates to a method of inhibiting TNF production in a mammal, including humans, which comprises administering to a mammal in need of such  
20 treatment, an effective TNF inhibiting amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. This method may be used for the prophylactic treatment or prevention of certain TNF mediated disease states effected thereby.

This invention further relates to a method of treating a human afflicted with a human immunodeficiency virus (HIV), which method comprises administering to a human  
25 in need of such treatment, an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

Compounds of Formula (I) are useful in the treatment of additional viral infections, where such viruses are sensitive to upregulation by TNF, or will elicit TNF production *in vivo*.

### 30 Detailed Description of the Invention

#### Definitions

As used herein, the following terms and expressions have the indicated meaning.

"Aryl" or "aralkyl", unless specified otherwise, means an aromatic ring or ring system of 6-10 carbon atoms, such as phenyl, benzyl, phenethyl, or naphthyl. The alkyl  
35 chain is meant to include both straight or branched chain radicals of 1 to 4 carbon atoms.

The term "C<sub>1-2</sub> alkyl", "C<sub>1-4</sub> alkyl", "C<sub>1-6</sub> alkyl" or "alkyl groups" includes both straight or branched chain radicals of 1 to 10 carbon atoms, unless the chain length is

otherwise limited thereto, including, but not limited to methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl, and the like.

The term "C<sub>3-7</sub> cycloalkyl" means groups of 3-7 carbon atoms, such as cyclopropyl, cyclopropylmethyl, cyclopentyl, or cyclohexyl.

5 "Cytokine" means any secreted polypeptide that affects the functions of cells, and is a molecule which modulates interactions between cells in immune, inflammatory, or hematopoietic responses. A cytokine includes, but is not limited to, monokines and lymphokines regardless of which cells produce them. The cytokine inhibited by the present invention for use in the treatment of an HIV-infected human must be a cytokine which is  
10 implicated in (a) the initiation and/or maintenance of T cell activation and/or activated T cell-mediated HIV gene expression and/or replication, and/or (b) any cytokine-mediated disease associated problem such as cachexia or muscle degeneration.

"Halo" includes all halogen radicals, i.e., chloro, fluoro, bromo, or iodo.

"Heteroaryl" means an aromatic ring system containing one or more heteroatoms,  
15 such as imidazolyl, triazolyl, oxazolyl, pyridyl, pyrimidyl, pyrazolyl, pyrrolyl, furanyl, or thienyl.

"Inhibiting the production of IL-1" or "inhibiting the production of TNF" means:

a) a decrease of excessive *in vivo* IL-1 or TNF levels in a human, to normal levels or below normal levels by inhibition of the *in vivo* release of IL-1 by all cells,  
20 including but not limited to monocytes or macrophages;

b) a down regulation, at the translational or transcriptional level, of excessive *in vivo* IL-1 or TNF levels in a human, to normal levels or below normal levels;  
or

c) a down regulation, by inhibition of the direct synthesis of IL-1 or TNF  
25 levels as a postranslational event.

"Percentage" and "%" refers to percentage by weight of a component or ingredient based on the weight of the total composition containing such component or ingredient.

"TNF mediated disease or disease states" means any and all disease states in which TNF plays a role, either by production of TNF itself, or by TNF causing another cytokine  
30 to be released, such as but not limited to IL-1 or IL-6. A disease state in which IL-1, for instance, is a major component, and whose production or action is exacerbated or secreted in response to TNF, would therefore be considered a disease state mediated by TNF. As TNF- $\beta$  (also known as lymphotoxin) has close structural homology with TNF- $\alpha$  (also known as cachectin), and since each induces similar biologic responses and binds to the  
35 same cellular receptor, both TNF- $\alpha$  and TNF- $\beta$  are inhibited by the compounds of the present invention and thus are herein referred to collectively as "TNF" unless specifically delineated otherwise.



This invention relates to a method for mediating or inhibiting the enzymatic activity or catalytic activity of PDE IV in a mammal in need thereof and for inhibiting the production of TNF in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

PDE IV inhibitors are useful in the treatment of a variety of allergic and inflammatory diseases, including: asthma, chronic bronchitis, atopic dermatitis, urticaria, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, eosinophilic granuloma, psoriasis, rheumatoid arthritis, septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, chronic glomerulonephritis, endotoxic shock and adult respiratory distress syndrome. In addition, PDE IV inhibitors are useful in the treatment of diabetes insipidus and central nervous system disorders such as depression and multi-infarct dementia.

The viruses contemplated for treatment herein are those that produce TNF as a result of infection, or those which are sensitive to inhibition, such as by decreased replication, directly or indirectly, by the TNF inhibitors of Formula (I). Such viruses include, but are not limited to HIV-1, HIV-2 and HIV-3, cytomegalovirus (CMV), influenza, adenovirus and the Herpes group of viruses, such as, but not limited to, *Herpes zoster* and *Herpes simplex*.

This invention more specifically relates to a method of treating a mammal, afflicted with a human immunodeficiency virus (HIV), which comprises administering to such mammal an effective TNF inhibiting amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

The compounds of this invention may also be used in association with the veterinary treatment of animals, other than humans, in need of inhibition of TNF production. TNF mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted above, but in particular viral infections. Examples of such viruses include, but are not limited to feline immunodeficiency virus (FIV) or other retroviral infection such as equine infectious anemia virus, caprine arthritis virus, visna virus, maedi virus and other lentiviruses.

The compounds of this invention are also useful in treating yeast and fungal infections, where such yeast and fungi are sensitive to upregulation by TNF or will elicit TNF production *in vivo*. A preferred disease state for treatment is fungal meningitis. Additionally, a compound of Formula (I) may be administered in conjunction with other drugs of choice for systemic yeast and fungal infections. Drugs of choice for fungal infections, include but are not limited to the class of compounds called the polymycins, such as Polymycin B, the class of compounds called the imidazoles, such as clotrimazole, econazole, miconazole, and ketoconazole; the class of compounds called the triazoles, such

as fluconazole, and itranazole, and the class of compounds called the Amphotericins, in particular Amphotericin B and liposomal Amphotericin B.

- 5 A compound of Formula (I) may also be used for inhibiting and/or reducing the toxicity of an anti-fungal, anti-bacterial or anti-viral agent by administering an effective amount of a compound of Formula (I) to a mammal in need of such treatment. Preferably, a compound of Formula (I) is administered for inhibiting or reducing the toxicity of the Amphotericin class of compounds, in particular Amphotericin B.


#### Preferred Compounds

- 10 The preferred compounds of this invention are those where  $R_1$  is OH or -COOH, X is  $YR_2$  particularly where Y is O and  $R_2$  is methyl substituted by 1 or more halogens;  $R_3$  is CN or  $-C\equiv CH$  and Z is a carboxylic acid derivative such as a  $-C(O)OH$ , a salt thereof or an ester or amide derivative such as  $C(=Y)R_{14}$ ,  $C(=O)OR_{14}$ ,  $C(=Y)NR_{10}R_{14}$ ,  $C(=NR_{10})NR_{10}R_{14}$ . In addition, it is preferred that the  $R_3$  group be axial and the Z group be equatorial.

- 15 The specific compounds disclosed herein are:  
 cis-{4-cyano-4-[3-(cis-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid};  
 cis-{4-cyano-4-[3-(cis-3-formyloxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid}, and  
 20 cis-{4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid}.

#### Syntheses

- Preparation of the compounds of Formula (I) can be carried out by one of skill in the art according to the procedures outlined reaction scheme set forth below and the specific chemistries set out in the Examples, infra. While the scheme and the examples illustrate the preparation of the cis/cis isomer(s), the cis/trans isomer(s) can be prepared by the same set of chemistries, with a nominal change in the treatment of compound 4; saponifying compound 4 provides the cis/trans compound namely the cis-{4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid} or the corresponding compounds as defined by Formula I. The preparation of any remaining  
 30 compounds of the Formula (I) not described therein may be prepared by the analogous processes disclosed herein which comprise:



5

## Example 1

Preparation of cis-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid]1(a) (3-Cyclopentyloxy-4-methoxyphenyl)acetonitrile

To a solution of 3-cyclopentyloxy-4-methoxybenzaldehyde (20 g, 90.8 mmol) in acetonitrile (100 mL) was added lithium bromide (15 g, 173 mmol) followed by the dropwise addition of trimethylsilylchloride (17.4 mL, 137 mmol). After 15 min, the reaction mixture was cooled to 0° C, 1,1,3,3-tetramethyldisiloxane (26.7 mL, 151 mmol) was added dropwise and the resulting mixture was allowed to warm to room temperature. After stirring for 3 h, the mixture was separated into two layers. The lower layer was

removed, diluted with methylene chloride and filtered through Celite®. The filtrate was concentrated under reduced pressure, dissolved in methylene chloride and refiltered. The solvent was removed in vacuo to provide a light tan oil. To a solution of this crude a-bromo-3-cyclopentyloxy-4-methoxytoluene in dimethylformamide (160 mL) under an argon atmosphere was added sodium cyanide (10.1 g, 206 mmol) and the resulting mixture was stirred at room temperature for 18 h, then poured into cold water (600 mL) and extracted three times with ether. The organic extract was washed three times with water, once with brine and was dried (K<sub>2</sub>CO<sub>3</sub>). The solvent was removed in vacuo and the residue was purified by flash chromatography (silica gel, 10% ethyl acetate/hexanes) to provide an off-white solid (m.p. 32-34° C); an additional quantity of slightly impure material also was isolated.

1(b) Dimethyl 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)pimelate

To a solution of (3-cyclopentyloxy-4-methoxyphenyl)acetonitrile (7 g, 30.3 mmol) in acetonitrile (200 mL) under an argon atmosphere was added a 40% solution of Triton-B in methanol (1.4 mL, 3.03 mmol) and the mixture was heated to reflux. Methyl acrylate (27 mL, 303 mmol) was added carefully, the reaction mixture was maintained at reflux for 5 h and then cooled. The mixture was diluted with ether, was washed once with 1N hydrochloric acid and once with brine, was dried (MgSO<sub>4</sub>) and the solvent was removed in vacuo. The solid residue was triturated with 5% ethanol/hexane to provide a white solid (m.p. 81-82° C); an additional quantity was also obtained from the filtrate. Anal. (C<sub>22</sub>H<sub>29</sub>NO<sub>6</sub>) calcd: C 65.49, H 7.25, N 3.47. found: C 65.47, H 7.11, N 3.49.

1(c) 2-Carbomethoxy-4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-one

To a suspension of sodium methoxide (350 mL, 1.55 mol, 25% w/w in methanol) in toluene (2.45 L) heated to 80° C under a nitrogen atmosphere was added a solution of dimethyl 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)pimelate (350.0 g, 0.87 mol) in toluene (1.05 L) over 10 min. The reaction was heated to 85° C by distilling away 250 mL of solvent and was vigorously stirred under nitrogen for 2 hours. The reaction was cooled to 50° C and was quenched with 3N (aq) HCl (700 mL, 2.1 mol). The organic layer was isolated, was washed once with deionized water (700 mL) and once with brine (700 mL). The organic layer was concentrated via low vacuum distillation to afford crude 2-carbomethoxy-4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-one in toluene. This was dissolved in 4.2 L of dimethyl sulfoxide and used in the next step.

1(d) 4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-one

To a suspension of sodium chloride (315 g, 5.39 mol) and deionized water (315 mL) was added the dimethyl sulfoxide (4.2 L) solution of 2-carbomethoxy-4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-one (323 g, 0.87 mol) and the resulting suspension was heated to 155° C for 1.75 h. The reaction was cooled to 40° C, was quenched into 8 L of iced water (2° C) and was extracted with ethyl acetate (3.5 L). The aqueous layer was isolated and re-extracted with 2.5 L of ethyl acetate. The combined organic extract (6 L) was washed two times with deionized water (2 x 1 L) and once with brine (1 L). The organic layer was isolated and concentrated in vacuo to afford a residue. This residue was dissolved in refluxing isopropanol (500 mL), was cooled to 0° C and held at this temperature for 1 hour. The crystals were isolated by filtration, were washed with 250 mL of isopropanol (0° C), and were dried in a vacuum oven (45° C at 20 inches) to produce 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-one. m.p. 111-112° C; Anal. (C<sub>19</sub>H<sub>23</sub>NO<sub>3</sub>) calcd: C 72.82, H 7.40, N 4.47; found: C 72.72, H 7.39, N 4.48.

1(e) 2-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexylidene]-1,3-dithiane

To a solution of 2-trimethylsilyl-1,3-dithiane (9.25 mL, 48.7 mmol) in dry tetrahydrofuran (80 mL) at 0° C under an argon atmosphere was added rapidly n-butyllithium (2.5M in hexanes, 19.2 mL, 48 mmol). After 10 min, the mixture was cooled to -78° C and a solution of 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-one (7.53 g, 23 mmol) in tetrahydrofuran (40 mL) was added. After 10 min, aqueous sodium chloride was added, the mixture was allowed to warm to room temperature and was diluted with water. This mixture was combined with the product of three substantially similar reactions conducted on ketone (3.04, 6.01 and 6.1 g, 48.3 mmol total), the combined mixture was extracted three times with methylene chloride, the extract was dried (MgSO<sub>4</sub>) and evaporated. Purification by flash chromatography (silica gel, 10% ethyl acetate/hexanes) provided a white solid. m.p. 115-116° C.

1(f) cis-[4-Cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid]

To a suspension of 2-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexylidene]-1,3-dithiane (140.0 g, 0.34 mol) in acetonitrile (500 mL) and deionized water (140 mL) under nitrogen was added trifluoroacetic acid (136 g, 1.19 mol). The suspension was heated to 65° C for 1.25 h followed by the addition of 20% sodium hydroxide (420 g, 2.1 mol). The solution was heated at 70 to 75° C for an additional 1.25 h, was cooled to 45° C, deionized water (420 mL) was added followed by 3N (aq) HCl (392 mL, 1.18 mol). The suspension was cooled to 5° C and held for 1 h. The suspension was filtered, was washed with cold (5° C) deionized water (200 mL), and was dried in a vacuum oven (40° C at 20 inches) to obtain crude cis-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid]. This material was assayed at 98.5% and was found to a 98.8:1.2 mixture of cis-to-trans isomers, which was contaminated with

0.1% of residual 1,3-propanedithiol. This material was purified via an oxidative workup as follows.

To a hot solution (65° C) of crude cis-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid] (85 g, 0.247 mol) in acetonitrile (425 mL) was added 1M sodium hydroxide (425 mL, 0.425 mol). To the solution (60° C) was added 4.25 g of calcium hypochlorite and the suspension was vigorously stirred for 2 h. The reaction was concentrated by distilling out 320 mL of solvent, followed by the addition of ethyl acetate (425 mL). The reaction was again concentrated by distilling out 445 mL of solvent, was cooled to 55° C followed by the addition of ethyl acetate (1.0 L) and 6N (aq.) HCl (100 mL). The organic layer was isolated, was washed three times with deionized water (3 x 300 mL), was filtered and was concentrated by distilling out 530 mL of solvent. To the solution was added ethyl acetate (635 mL) with continued distillation to remove 750 mL of solvent. The solution was cooled to 65° C followed by the addition of hexane (340 mL). The suspension was cooled to 5° C, held at this temperature for 1 hour, was filtered and was washed with cold (5° C) 10% ethyl acetate/hexane (200 mL). The solid was collected and was dried in a vacuum oven (40° C at 20 inches) to obtain cis-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid]. This material was found to contain no trans isomer. Anal. (C<sub>20</sub>H<sub>25</sub>NO<sub>4</sub>) calcd: C 69.95, H 7.34, N 4.08; found: C 69.90, H 7.35, N 4.02.

20

#### Example 2

##### Preparation of cis-[4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]-cyclohexane-1-carboxylic acid]

##### 2(a) cis-[4-Cyano-4-(3-hydroxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid]

To a solution of boron tribromide in dichloromethane (0.1M, 335 mL, 33.5 mmol) under an argon atmosphere at -78° C was slowly added a solution of cis-[4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexane-1-carboxylic acid] (4.03 g, 11.7 mmol) in dichloromethane (180 mL). The mixture was stirred for 5 min, 15% sodium methoxide in methanol was added to pH 8-9 and the reaction was warmed to RT. Water (100mL) was added and the mixture was acidified with 3N aqueous hydrochloric acid to pH 1-2. The organic layer was separated, was dried (MgSO<sub>4</sub>/Na<sub>2</sub>SO<sub>4</sub>), was filtered and was evaporated. The residue was twice dissolved in chloroform and the solution was evaporated to yield a white solid. <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>) δ 7.01 (d, J=2.4 Hz, 1H), 6.96 (d of d, J=2.4, 8.5 Hz, 1H), 3.89 (s, 3H), 2.31 (m, 1H), 2.21 (br t, J=13.6 Hz, 4H), 1.98 (m, 2H), 1.77 (m, 2H); mp 190-193° C.

35

##### 2(b) Methyl cis-[4-cyano-4-(3-hydroxy-4-methoxyphenyl)cyclohexane-1-carboxylate]

*p*-Toluenesulfonic acid monohydrate (0.015 g, 0.08 mmol) was added to a solution of the compound of Example 2(a) (0.70 g, 2.54 mmol) in dry methanol (20 mL) under an argon atmosphere and the reaction was stirred for 6 h at 45-50° C. The reaction was

cooled to RT and was stirred for an additional 16 h. The solution was evaporated and the residue was purified by flash chromatography (silica gel, 50% hexane/ethyl acetate) to yield the title compound as a white solid. <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>) δ 7.01 (m, 2H), 6.85 (d, J=9.1 Hz, 1H), 3.90 (s, 3H), 3.72 (s, 3H), 2.35 (t of t, J=3.6, 12.2 Hz, 1H), 2.14-2.25 (m, 4H), 2.00 (app q, J=13.4 Hz, 1H), 1.99 (app q, J=13.4 Hz, 1H), 1.77 (app t, J=13.4 Hz, 1H), 1.76 (app t, J=13.4 Hz, 1H); mp 106-107° C.

2(c) Methyl cis-(4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]-cyclohexane-1-carboxylate)

The compound of Example 2(b) (0.69 g, 2.37 mmol) was dissolved in tetrahydrofuran (20 mL) under an argon atmosphere and was treated with triphenylphosphine (1.24 g, 4.74 mmol) and cis-1,3-cyclopentanediol (0.49 g, 4.74 mmol). Diethyl azodicarboxylate (0.83 g, 4.74 mmol) was added and the mixture was stirred at RT for 16 h. The solution was evaporated, the residue was diluted with ether and the white solid was removed by filtration. The filtrate was concentrated and the residue was purified by flash chromatography (silica gel, 50% hexane/ethyl acetate) to yield a mixture of the title compound and triphenylphosphine oxide. The mixture was diluted with ether and the white solid triphenylphosphine oxide was removed by filtration. Evaporation of the filtrate yielded the title compound as a sticky, colorless semi-solid. <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>) δ 7.07 (d, J=2.4 Hz, 1H), 7.02 (d of d, J=2.4, 8.8 Hz, 1H), 6.87 (d, J=8.8 Hz, 1H), 4.99 (m, 1H), 4.37 (m, 1H), 3.85 (s, 3H), 3.74 (s, 3H), 3.16 (d, J=9.1 Hz, 1H), 2.39 (m, 1H), 1.88-2.25 (m, 12H), 1.80 (br t, J=13.5 Hz, 2H).

2(d) cis-(4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid)

The compound of Example 2(c) (0.10 g, 0.27 mmol) was dissolved in 5:5:2 tetrahydrofuran/methanol/water (5 mL), sodium hydroxide (0.035 g, 0.88 mmol) was added and the mixture was stirred at RT for 3 h. The solvent was evaporated, the residue was partitioned between 5% aqueous NaOH and dichloromethane and the layers were separated. The aqueous layer was acidified to pH 3 with 3N aqueous hydrochloric acid and was extracted three times with 5% methanol in chloroform. The organic extracts were combined, were dried (MgSO<sub>4</sub>), filtered and evaporated. The residue was purified by flash chromatography (silica gel, 90:10:1 chloroform/methanol/water) to yield a solid which was slurried in ether, was collected by filtration and was dried in vacuo to afford the title compound. MS(Cl/NH<sub>3</sub>) m/e 377 [M + NH<sub>3</sub>]<sup>+</sup>; <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>) δ 7.08 (br s, 1H), 7.03 (br d, J=8.5 Hz, 1H), 6.88 (d, J=8.5 Hz, 1H), 4.98 (m, 1H), 4.38 (m, 1H), 3.84 (s, 1H), 2.41 (m, 1H), 1.77-2.29 (m, 16H); Anal. (C<sub>20</sub>H<sub>25</sub>NO<sub>5</sub>•0.9 H<sub>2</sub>O) calcd: C, 63.95; H, 7.19; N, 3.73. found: C, 64.06; H, 6.88; N, 3.77; mp 161-163° C.

Example 3

Preparation of cis-[4-cyano-4-[3-(cis-3-hydroxycyclopentyloxy)-4-methoxyphenyl]-cyclohexane-1-carboxylic acid]

3(a) Methyl cis-[4-cyano-4-[3-(cis-3-formyloxycyclopentyloxy)-4-methoxyphenyl]-cyclohexane-1-carboxylate]

5       The compound of Example 2(c) (0.68 g, 1.83 mmol) was dissolved in tetrahydrofuran (20 mL) under an argon atmosphere and was treated with triphenylphosphine (0.96 g, 3.66 mmol) and formic acid (0.17 g, 3.66 mmol). Diethyl azodicarboxylate (0.64 g, 3.66 mmol) was added and the mixture was stirred at RT for 16 h. The solution was evaporated, ether was added and the white solid was removed by  
10       filtration. The filtrate was concentrated and the residue was purified by flash chromatography (silica gel, 65% hexane/ethyl acetate) to yield the title compound as a clear colorless oil. <sup>1</sup>H NMR(400 MHz, CDCl<sub>3</sub>) δ 8.02 (s, 1H), 7.0 (d of d, J=2.4, 8.2 Hz, 1H), 6.99 (d, J=2.4 Hz, 1H), 6.87 (d, J=8.2 Hz, 1H), 5.48 (m, 1H), 4.95 (m, 1H), 3.84 (s, 3H), 3.72 (s, 3H), 2.31-2.40 (m, 2H), 2.13-2.28 (m, 7H), 1.96-2.06 (m, 3H), 1.74-1.87 (m,  
15       3H).

3(b) cis-[4-cyano-4-[3-(cis-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid]

      The compound of Example 3(a) (0.52 g, 1.31 mmol) was dissolved in 5:5:2 tetrahydrofuran/methanol/water (20mL), sodium hydroxide (0.32 g, 8.0 mmol) was added  
20       and the mixture was stirred at RT for 2.5 h. The solvent was evaporated and the aqueous residue was acidified to pH 1-2 with 3N aqueous hydrochloric acid. The white solid product was collected, was washed with water and was dried in vacuo to afford the title compound as a white solid. MS(CI/NH<sub>3</sub>) m/e 377 [M + NH<sub>3</sub>]<sup>+</sup>; <sup>1</sup>H NMR(250 MHz, CDCl<sub>3</sub>) δ 6.98 (m, 2H), 6.86 (d, J=8.2 Hz, 1H), 4.97 (m, 1H), 4.59 (m, 1H), 3.85 (s, 3H),  
25       1.64-2.47 (m, 17H); mp 143-145° C.

**METHODS OF TREATMENT**

      In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof may be used neat though a preferred technique is to present them with a carrier/diluent accordance with standard pharmaceutical practice. Any formulation  
30       compatible with the chosen method of delivery and the stability of the compound may be used. One skilled in the art will be able to select and prepare an acceptable formulation in accordance with standard practices in the field of the formulary arts.

      The compounds of Formula (I) or may be administered orally (when active by this route), oral, intravenous, intraperitoneal, and intramuscular administration, topically,  
35       parenterally, or by inhalation in conventional dosage forms prepared by combining such agent with standard pharmaceutical carriers according to conventional procedures in an amount sufficient to produce the desired therapeutic activity.



The amount of a compound of Formula (I) required for therapeutic effect on topical administration will, of course, vary with the compound chosen, the nature and severity of the condition and the animal undergoing treatment, and is ultimately at the discretion of the physician.

- 5        The daily dosage regimen for oral administration is suitably about .001 mg/kg to 100mg/kg, preferably 0.01 mg/Kg to 40 mg/Kg, of a compound of Formula (I) or a pharmaceutically acceptable salt thereof calculated as the free base. The active ingredient may be administered from 1 to 6 times a day, sufficient to exhibit activity.

#### UTILITY EXAMPLES

10

##### Example A

##### Inhibitory effect of compounds of Formula (I) on *in vitro* TNF production by human monocytes

- 15        The inhibitory effect of compounds of Formula (I) on *in vitro* TNF production by human monocytes may be determined by the protocol as described in Badger *et al.*, EPO published Application 0 411 754 A2, February 6, 1991, and in Hanna, WO 90/15534, December 27, 1990.

##### Example B

- 20        Two models of endotoxic shock have been utilized to determine *in vivo* TNF activity for the compounds of Formula (I). The protocol used in these models is described in Badger *et al.*, EPO published Application 0 411 754 A2, February 6, 1991, and in Hanna, WO 90/15534, December 27, 1990.

The exemplified compounds herein demonstrated a positive *in vivo* response in reducing serum levels of TNF induced by the injection of endotoxin.

##### Example C

- 25        Isolation of PDE Isozymes

- 30        The phosphodiesterase inhibitory activity and selectivity of the compounds of Formula (I) can be determined using a battery of five distinct PDE isozymes. The tissues used as sources of the different isozymes are as follows: 1) PDE Ib, porcine aorta; 2) PDE Ic, guinea-pig heart; 3) PDE III, guinea-pig heart; 4) PDE IV, human monocyte; and 5) PDE V (also called "Ia"), canine trachealis. PDEs Ia, Ib, Ic and III are partially purified using standard chromatographic techniques [Torphy and Cieslinski, Mol. Pharmacol., 37:206-214, 1990]. PDE IV is purified to kinetic homogeneity by the sequential use of anion-exchange followed by heparin-Sepharose chromatography [Torphy *et al.*, J. Biol. Chem., 267:1798-1804, 1992].

- 35        Phosphodiesterase activity is assayed as described in the protocol of Torphy and Cieslinski, Mol. Pharmacol., 37:206-214, 1990. Positive IC<sub>50</sub>'s in the nanomolar to  $\mu$ M range for compounds of the workings examples described herein for Formula (I) have been demonstrated.


Example D

The ability of selected PDE IV inhibitors to increase cAMP accumulation in intact tissues is assessed using U-937 cells, a human monocyte cell line that has been shown to contain a large amount of PDE IV. To assess the activity of PDE IV inhibition in intact  
5 cells, nondifferentiated U-937 cells (approximately  $10^5$  cells/reaction tube) were incubated with various concentrations (0.01-1000  $\mu$ M) of PDE inhibitors for one minute and 1 $\mu$ M prostaglandin E2 for an additional four minutes. Five minutes after initiating the reaction, cells were lysed by the addition of 17.5% perchloric acid, the pH was neutralized by the addition of 1M potassium carbonate and cAMP content was assessed by RIA. A  
10 general protocol for this assay is described in Brooker *et al.*, Radioimmunoassay of cyclic AMP and cyclic GMP., Adv. Cyclic Nucleotide Res., 10:1-33, 1979. The compounds of the working examples as described herein for Formula (I) have demonstrated a positive EC<sub>50</sub>s in the  $\mu$ M range in the above assay.

No toxic effects are expected when these compounds are administered in  
15 accordance with the present invention.

What is claimed is:

1. A compound of Formula I



5 wherein

$R_1$  is OH or an ether or ester thereof,

X is  $YR_2$ , halogen, nitro,  $NR_4R_5$  or formyl amine;

Y is O or  $S(O)_m$ , where m is 0, 1 or 2;

10  $R_2$  is methyl or ethyl, where either methyl or ethyl may be optionally substituted by 1 or more halogens;

$R_3$  is hydrogen, halogen,  $C_{1-4}$ alkyl,  $CH_2NHC(O)C(O)NH_2$ , halo-substituted  $C_{1-4}$ alkyl,  $-CH=CHR_8 \cdot R_8'$ , cyclopropyl optionally substituted by  $R_8'$ , CN,  $OR_8$ ,  $CH_2OR_8$ ,  $NR_8R_{10}$ ,  $CH_2NR_8R_{10}$ ,  $C(Z')H$ ,  $C(O)OR_8$ ,  $C(O)NR_8R_{10}$ , or  $-C \equiv CR_8$ ;

$R_4$  and  $R_5$  are independently hydrogen or  $C_{1-2}$  alkyl;

15  $R_7$  is  $-(CR_4R_5)_qR_{12}$  or  $C_{1-6}$  alkyl wherein the  $R_{12}$  or  $C_{1-6}$  alkyl group is optionally substituted one or more times by  $C_{1-2}$  alkyl optionally substituted by one to three groups selected from -F, -Br, -Cl,  $-NO_2$ ,  $-NR_{10}R_{11}$ ,  $-C(=O)R_8$ ,  $-C(=O)OR_8$ ,  $-OR_8$ , -CN,  $-C(=O)NR_{10}R_{11}$ ,  $-OC(=O)NR_{10}R_{11}$ ,  $-OC(=O)R_8$ ,  $-NR_{10}C(=O)NR_{10}R_{11}$ ,  $-NR_{10}C(=O)R_{11}$ ,  $-NR_{10}C(=O)OR_9$ ,  $-NR_{10}C(=O)R_{13}$ ,  $-C(=NR_{10})NR_{10}R_{11}$ ,  $-C(=N-$   
 20  $CN)NR_{10}R_{11}$ ,  $-C(=N-CN)SR_9$ ,  $-NR_{10}C(=N-CN)NR_{10}R_{11}$ ,  $-NR_{10}S(=O)_2R_9$ ,  $-S(O)_mR_9$ ,  $-NR_{10}C(=O)C(=O)NR_{10}R_{11}$ ,  $-NR_{10}C(=O)C(=O)R_{10}$ , thiazolyl, imidazolyl, oxazolyl, pyrazolyl, triazolyl or tetrazolyl;

$R_8$  is -H or  $R_9$ ;

$R_8'$  is  $R_8$  or fluorine;

25  $R_9$  is  $C_{1-4}$  alkyl optionally substituted by one to three -F;

$R_{10}$  is  $OR_8$ , hydrogen, or  $C_{1-4}$ alkyl optionally substituted by one to three fluorines;

$R_{11}$  is -H or  $C_{1-4}$  alkyl optionally substituted by one to three -F; or when  $R_{10}$  and  $R_{11}$  are as  $NR_{10}R_{11}$  they may together with the nitrogen form a 5 to 7 membered ring  
 30 optionally containing at least one additional heteroatom selected from O, N or S;

$R_{12}$  is  $C_{3-7}$  cycloalkyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrimidyl, pyrazolyl, 1-imidazolyl, 2-imidazolyl, thiazolyl, triazolyl, pyrrolyl, piperazinyl, piperidinyl, morpholinyl, furanyl, 2-thienyl, 3-thienyl, 4-thiazolyl, 5-thiazolyl, quinolinyl, naphthyl or phenyl;

$R_{13}$  is a heterocyclic ring selected from oxazolidinyl, oxazolyl, thiazolyl, pyrazolyl, triazolyl, tetrazolyl, imidazolyl, imidazolidinyl, thiazolidinyl, isoxazolyl, oxadiazolyl or thiadiazolyl, where  $R_{13}$  is appended to a compound of Formula (I) through a carbon atom of the heterocyclic ring, and where each heterocyclic ring may be unsubstituted or substituted by one or two  $C_{1-2}$  alkyl groups;

$R_{14}$  is H or  $R_7$ , or when  $R_{10}$  and  $R_{14}$  are as  $NR_{10}R_{14}$ , they may together with the nitrogen atom form a 5 to 7 membered ring optionally containing at least one additional heteroatom selected from O, N or S;

$m'$  is 0, 1 or 2;

$q$  is 0, 1 or 2.

$Y'$  is O or S;

$Z$  is  $C(=Y')R_{14}$ ,  $C(=O)OR_{14}$ ,  $C(=Y')NR_{10}R_{14}$ ,  $C(=NR_{10})NR_{10}R_{14}$ , CN,  $C(=NOR_8)R_{14}$ ,  $C(=O)NR_8NR_8C(=O)R_8$ ,  $C(=O)NR_8NR_{10}R_{14}$ ,  $C(=NOR_{14})R_8$ ,  $C(=NR_8)NR_{10}R_{14}$ ,  $C(=NR_{14})NR_8R_8$ ,  $C(=N-CN)NR_{10}R_{14}$ ,  $C(=N-CN)SR_9$ , 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 4-triazolyl[1,2,3], 5-triazolyl[1,2,3], 3-triazolyl[1,2,4], 5-triazolyl[1,2,4], 5-tetrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-oxadiazolyl[1,2,4], 5-oxadiazolyl[1,2,4], 2-oxadiazolyl[1,3,4], 2-thiadiazolyl[1,3,4], 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-oxazolidinyl, 4-oxazolidinyl, 5-oxazolidinyl, 2-thiazolidinyl, 4-thiazolidinyl or 5-thiazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, or 5-imidazolidinyl; wherein all of the heterocyclic ring systems may be optionally substituted one or more times by  $R_{14}$ ;

$Z'$  is O,  $NR_9$ ,  $NOR_8$ ,  $NCN$ ,  $C(-CN)_2$ ,  $CR_8CN$ ,  $CR_8NO_2$ ,  $CR_8C(O)OR_8$ ,  $CR_8C(O)NR_8R_8$ ,  $C(-CN)NO_2$ ,  $C(-CN)C(O)OR_9$ , or  $C(-CN)C(O)NR_8R_8$ ;

or

a pharmaceutically acceptable salt thereof.

2. A compound of claim 1 wherein  $R_1$  is OH or  $-COOH$ ,  $X$  is  $YR_2$ ,  $Y$  is O,  $R_2$  is methyl substituted by 1 or more halogens;  $R_3$  is CN or  $-C\equiv CH$ ,  $Z$  is a carboxylic acid derivative such as a  $-C(O)OH$ , a salt thereof or an ester or amide derivative such as  $C(=Y')R_{14}$ ,  $C(=O)OR_{14}$ ,  $C(=Y')NR_{10}R_{14}$ ,  $C(=NR_{10})NR_{10}R_{14}$  and the  $R_3$  group is axial and the  $Z$  group is equatorial.

3. A compound of claim 2 which is


cis-{4-cyano-4-[3-(cis-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid};

cis-{4-cyano-4-[3-(cis-3-formyloxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid}, or

cis-{4-cyano-4-[3-(trans-3-hydroxycyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid}, or a pharmaceutically acceptable salt thereof.


4. A method for treating asthma comprising a compound according to any one of claims 1-3 and a pharmaceutically acceptable excipient.
5. A pharmaceutical composition comprising a compound of according to any one of claims 1-3.
6. A process for preparing the ketone of formula II

10




where R<sub>1</sub> is hydrogen or another a substituent, which process comprises cyclizing a compound of formula A

15




where alk is alkyl of 1 to 6 carbon atoms by mixing the ester with a strong base in a non-polar solvent and heating the mixture to between 50 and 100° C under an inert atmosphere for a period of between 30 minutes and 4 hours.

7. The process of claim 6 wherein the base is an alkali metal alkoxide.
8. The process of any one of claims 6 or 7 wherein the base is sodium methoxide and the solvent is toluene.
9. The process of any one of claims 6 to 8 wherein the product is 2-carbomethoxy-4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-one.
10. A process for preparing an acid of formula IV



wherein  $R_1$  is hydrogen or another substituent and X is OH or a salt thereof or  $-O-C_1-$ alkyl, which process comprises hydrolyzing a thioketene of formula (C)

5



using an acid and aqueous solvent system and heating the mixture to between 40 and 80° C for 30 minutes to 2 hours, and then adding a base.

10

11. The process of claim 10 wherein the acid is an organic acid.

12. The process of either claim 10 or 11 wherein the acid is trifluoroacetic acid.

13. The process of any one of claims 10 to 12 wherein the product is an acid of formula IV which is cis-[4-cyano-4-(3-cyclopentyl-4-methoxyphenyl)cyclohexane-1-carboxylic acid] or a salt thereof.

15

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/02965**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C07C 253/30, 255/36, 255/37, 255/38, 255/41, 255/46; A61K 31/275

US CL : 514/521; 558/365, 426

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/521; 558/365, 426

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C.A.S. online

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstracts, Volume 120, issued 1994, Christensen, "Cyclohexylbenzene derivatives useful for treating allergic and inflammatory diseases", Abstract No. 120: 134020m, Abstract of PCT Int. Appl. WO 93/19,747, (14 October 1993), page 29.	1-8, 10-12
A	Chemical Abstracts, Volume 120, issued 1994, Christensen, et al. "Preparation of arylcyclohexanes useful and for inhibiting production of tumor necrosis factor" Abstract No. 120: 163532k, Abstract of PCT Int. Appl. WO 93/19,748, (14 October 1993), page 27.	1-8, 10-12

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

07 JUNE 1995

Date of mailing of the international search report

30 JUN 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOSEPH PAUL BRUST

Telephone No. (703) 308-1235

**INTERNATIONAL SEARCH REPORT**International application No.  
PCT/US95/02965**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstracts, Volume 120, issued 1994 Christensen, et al., "Cyclohexylbenzones useful for treating allergic or inflammatory diseases", Abstract No. 120: 191326q, Abstract of PCT Int. Appl. WO 93/19,750, (14 October 1993), page 56.	1-8, 10-12



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/02965

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 1, 2, 4, 5, 6-8, 10-12 (all in part)  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
Please See Extra Sheet.
  
3. ☒ Claims Nos.: 9 and 13  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/02965

## BOX I. OBSERVATIONS WHERE CLAIMS WERE FOUND UNSEARCHABLE

### 2. Where no meaningful search could be carried out, specifically:

Concerning compound, composition and method of use claims 1, 2, 4 and 5, aside from the three specific structures named in claim 3 (and the corresponding disclosure of these three compounds in the specification) the terms otherwise used in these unsearched claims 1, 2, 4 and 5 cannot be reasonably or directly ascertained into any meaningful enough specific compound structures so as to afford a reasonable determination of proper specific subclasses to search. Thus, these unsearchable claims 1, 2, 4 and 5 will be searched to the extent that they read on searchable features (i.e., all compounds of claim 3 with the subject matter area being defined as non-heterocyclic group containing compounds but containing a 4-cyano-4-[-3-(oxycyclopentyloxy)-4-alkoxyphenyl] cyclohexane-1-carboxylic acids "core").

Concerning chemical process for preparing compounds of formula II, claims 6-8 and chemical process for preparing compounds of formula IV, claims 10-12, the same cannot be reasonably ascertained into any meaningful enough specific prepared products. This is due to the R1 being defined as "another a substituent" in the product of claims 6-8 and as "another substituent" in the product of claims 10-12 and the product of claims 10-12 being peroxidic in nature (i.e., the product of formula IV containing a moiety "-COOX" where "X is OH" or "-O-C1-6 alkyl" which is ultimately — -COOOH or -(COOO-C6 alkyl) which is not the product of the 1(e) to 1(f) reaction on page 11, lines 15 to 37. The search can, and has, been carried out on the preparation of compounds of formula II in claims 6-8 and the preparation of compounds of formula IV in claims 10-12 all wherein the R1's are H and/or alkoxy and/or oxy substituted cycloalkoxy and wherein the -COOX of formula IV is interpreted as -COOH or -COO-C1-6 alkyl.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**